



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The condition of plague in the south remains about the same. During the week 10 cases and 2 deaths have been reported from Payta. I received information from a reliable source that yellow fever is present in Buenaventura. As this is the last port at which the coasting steamers stop before reaching Ancon and is only about thirty-five hours distant, I cabled Surg. H. R. Carter, chief quarantine officer Isthmian Canal Commission.

Report for month of July, 1904—Grippe epidemic.

Week ended August 3, 1904, mortality from all causes, 53, as follows: Yellow fever, 3; pernicious fever, 4; fever without classification, 6; grippe, 3; tuberculosis, 9; enteric diseases, 12; from all other causes, 16. During the month ended July 31 there were 223 deaths from all causes, of which 24 were from yellow fever and 27 from tuberculosis. For the corresponding period last year there were 246 deaths, of which none were from yellow fever and 30 from tuberculosis. On August 2, a supplemental bill of health was issued to steamship *Santiago*, en route from Chilean and Peruvian ports to Ancon, C. Z. This vessel took 8 cabin and 10 steerage passengers from this port. Twenty-two cabin passengers from Callao, who wished to disembark here, were transferred to a small steamer and sent down the river to complete fifteen days' quarantine. One immune certificate was issued for Ancon and 36 pieces of baggage were inspected and labeled. This is in accordance with instructions received from Surg. H. R. Carter, of the Service, Chief Quarantine Officer Isthmian Canal Commission. Grippe, which on June 24 I reported as prevailing, has assumed the proportion of a mild epidemic.

Rats dying on island of Puná—Suspect plague.

On August 5 the medical officer of the sanitary station of Puná reported to the superior board of health that rats were dying there and presented symptoms of plague. A meeting of the board was immediately called, and a committee appointed to go there and investigate. Simultaneously with the report from the medical officer at Puná were rumors of suspicious deaths there. Puná is a small island situated about 36 miles from Guayaquil, at the mouth of the river. At this writing no further information can be obtained.

GERMANY.

Report from Berlin—Plague and cholera in various countries.

Consul-General Mason reports, August 8, 1904, as follows:

Plague.

Egypt.—During the week ended July 23 there were registered 16 new cases of plague (and 8 deaths), namely, 6 cases (2 deaths) in Zagazig, 3 cases (3 deaths) in the district of Bibe, 3 cases (1 death) in Alexandria, 2 fresh cases in Samallut, 1 case (1 death) in Beni-Souef, and 1 case (1 death) in Port Said.